

On Group bijections ϕ with $\phi(B) = A$ and $\forall a \in B, a\phi(a) \notin A$

Y. O. Hamidoune*

December 13, 2008

Abstract

A *Wakeford pairing* from S onto T is a bijection $\phi : S \rightarrow T$ such that $x\phi(x) \notin T$, for every $x \in S$. The number of such pairings will be denoted by $\mu(S, T)$.

Let A and B be finite subsets of a group G with $1 \notin B$ and $|A| = |B|$. Also assume that the order of every element of B is $\geq |B|$.

Extending results due to Losonczy and Eliahou-Lecouvey, we show that $\mu(B, A) \neq 0$.

Moreover we show that $\mu(B, A) \geq \min\{\frac{|B|+1}{3}, \frac{|B|(q-|B|-1)}{2q-|B|-4}\}$, unless there is $a \in A$ such that $|Aa^{-1} \cap B| = |B| - 1$ or Aa^{-1} is a progression.

In particular, either $\mu(B, B) \geq \min\{\frac{|B|+1}{3}, \frac{|B|(q-|B|-1)}{2q-|B|-4}\}$, or for some $a \in B$, Ba^{-1} is a progression.

MSC Classification: 11B60, 11B34, 20D60.

1 Introduction

Let A and B be finite subsets of a group G with $|A| = |B|$. A *Wakeford pairing* from B onto A is a bijection $\phi : B \rightarrow A$ such that $x\phi(x) \notin A$, for every $x \in B$.

Our pairings are dual to pairings used in litterature [2, 16, 1]. The two notions are equivalent up to replacing the group its opposite group or by

*UPMC Univ Paris 06, E. Combinatoire, Case 189, 4 Place Jussieu, 75005 Paris, France.

replacing (A, B) by (A^{-1}, B^{-1}) . With our choice, isoperimetric theorems apply more quicquely. Fan and Losonczy [2] introduced this notion in $\mathbb{Z}^n$ in connection with an old problem of Wakeford related to canonical forms for symmetric tensors.

The number of distinct matching from B onto A will be denoted by $\mu(B, A)$.

Let us define a *prime* group as a group having no proper finite subgroup H . By elementary Group Theory, a group is a prime group if and only if it is a torsion free group or if it has a prime order.

The next two results are due Losonczy in the abelian case [16] and to Eliahou-Lecouvey in the non-abelian case [1]:

- If A and B be finite subsets of a prime group with the same cardinality such that $1 \notin B$, then $\mu(B, A) \neq 0$.
- If B is a finite subset of a group such that $1 \notin B$, then $\mu(B, B) \neq 0$.

The relation $1 \notin B$ is obviously a necessary condition for the existence of such a pairing. Take a proper finite subgroup H , $h \in H \setminus \{1\}$ and $a \notin H$. Put $B = (H \setminus \{1\}) \cup \{a\}$. Suppose that there exists a Wakeford pairing $\phi : B \rightarrow H$. Clearly $\phi(a) = 1$ and hence $\phi(h) \in H$. Thus $\phi(h)h \in H$, a contradiction.

These observations made by Losonczy in [16] show that in a non-prime group, there exist finite subsets A and B with the same cardinality such that $1 \notin B$, and $\mu(B, A) = 0$. such a thing can not hold in prime groups, by the result mentioned above.

The results mentioned above are proved using some standard Addition Theorems presented below:

Let A and B be finite subsets of a group G . Kneser's Theorem [14] states that $|AB| \geq |A| + |B| - 1$ if G is abelian and if AB is aperiodic. The Scherck-Kemperman Theorem states that $|AB| \geq |A| + |B| - 1$, if $A \cap B^{-1} = \{1\}$. This result was proved first by Scherck [22] for abelian groups and by Kemperman [13] for arbitrary groups. As observed by Eliahou-Lecouvey [1], the existence of a symmetric pairing is related the Scherck-Kemperman Theorem. In the non-abelian case, a result due to Olson [21] states that $|AB| \geq |A| + |B| - |H|$, where H is a subgroup depending on A and B . This last result is related to one result proved independently and shortly before it by the author in [6]. Developments of this the last result are known as isoperimetric results:

A main tool in the present work is the isoperimetric approach developed by the author [6, 7, 8, 9, 11]. Let us present briefly this method:

The subgroup generated by a set X will be denoted by $\langle X \rangle$. Let S be a finite subset of a group G with $1 \in S$. The k th-connectivity of S is defined as

$$\kappa_k(S) = \min\{|\langle XS \rangle \setminus X| : \infty > |X| \geq k, X \subset \langle S \rangle \text{ and } |XS| \leq |\langle S \rangle| - k\},$$

where $\min \emptyset = |\langle S \rangle| - k + 1$.

We shall say that S is a *Cauchy subset* if $\kappa_1(S) = |S| - 1$. We shall say that S is a *Vosper subset* if $\kappa_2(S) \geq |S|$. Clearly S is a Cauchy subset if and only if for every $X \subset \langle S \rangle$ with $|X| \geq 1$,

$$|XS| \geq \min(|\langle S \rangle|, |X| + |S| - 1).$$

Also, S is a Vosper subset if and only if for every $X \subset \langle S \rangle$ with $|X| \geq 2$,

$$|XS| \geq \min(|\langle S \rangle| - 1, |X| + |S|).$$

A non-empty subset S of a group will be called a Chowla subset if the order of every element of S is $\geq |S| + 1$. The notion of a Chowla subset, introduced by the author in [8] as a relaxation of Chowla's condition in cyclic groups, allows to extend additive properties of prime groups to a large class of subsets of an arbitrary group.

Pairings existence is related to the Cauchy property and the Vosper's property allows to give a lower bound for the number of distinct pairings. We shall investigate pairing of a Chowla subset B onto an arbitrary A and obtain a lower bound for the number of distinct pairings in this case.

Recall the following notion used by Károlyi in his generalization of Vosper's Theorem [12]:

For a group G , put

$$p(G) = \min\{|M| : M \text{ is a finite subgroup of } G, \text{ with } 2 \leq |M| < |G|\},$$

where $\min \emptyset = \infty$. In particular $p(G) = \infty$ if G is a prime group.

The organization of the paper is the following:

Section 2 contains essentially known results. Section 2.1 presents König-Hall's Theorem and its particular formulation in the Wakeford graph. Section 2.2 the Erdős-Heilbronn averaging argument used to give a bound for

the maximum degree of the Wakeford graph $\mathcal{R} = \{(x, y) \in B \times A \mid xy \notin A\}$. Section 2.3 presents some isoperimetric formalism. In Section 3, we prove an inverse theorem for cofinite sets. As an application, we obtain the following result:

If B be a finite Chowla subset of a group G and A is a subset of G with $|A| = |B|$, then $\mu(B, A) \neq 0$.

In particular, $\mu(B, A) \neq 0$, if $|B| < p(G)$, where B is a finite subset of a group $G \setminus \{1\}$ and A is a subset of G with $|A| = |B|$.

Applied to groups with a prime order, the last result reduces to a result of Losonczy [16]. Also in the case of torsion free groups, it reduces to a result of Eliahou-Lecouvey [1] generalizing results of Wakeford [24] and Losonczy [16] in the abelian case [24].

Assuming that B is a finite Chowla subset of a group G and that A be a subset of G with $|A| = |B|$, we show in Section 4 that one of the following holds:

- (i) $\mu(B, A) \geq \min\{\frac{|B|+1}{3}, \frac{|B|(q-|B|-1)}{2q-|B|-4}\}$,
- (ii) For some $a \in A$, $|Aa^{-1} \cap B| = |B| - 1$,
- (iii) For some $a \in A$, Aa^{-1} is a progression,

where q denotes the cardinality of the subgroup generated by B . In particular either $\mu(B, B) \geq \min\{\frac{|B|+1}{3}, \frac{|B|(q-|B|-1)}{2q-|B|-4}\}$, or for some $a \in B$, Ba^{-1} is a progression.

Let us now construct two basic examples having a unique Wakeford pairing, where exactly one of the conditions (ii) and (iii) is satisfied:

Put $P = \{1, r, \dots, r^j\}$. There is clearly a unique matching from rP onto P , where $\phi : r^s \mapsto r^{j+1-s}$. Thus $\mu(rP, P) = 1$.

Put $P = \{1, r^2, \dots, r^{j+1}\}$ and take $a \notin P \cup \{r^{j+1}, r^{j+2}\}$. Set $Q = \{r^2, \dots, r^{j+1}, a\}$. There is a unique matching $\phi : Q \rightarrow P$ with $\phi(a) = 1$. Thus $\mu(Q, P) = 1$.

The last example shows that one may have $\mu(A, B) = 1$ for a set A which is not a right progression.

2 Preliminaries

2.1 Matchings

Let $\mathcal{R} \subset V \times W$ be a relation. A *selection* of $\mathcal{R}$ is a mapping $\mathcal{S} : V \rightarrow W$ such that $(x, \mathcal{S}(x)) \in \mathcal{R}$, for every $x \in V$. We shall write

$$\Delta(\mathcal{R}) = \max\{|\mathcal{R}(x)| : x \in V\} \text{ and } \Delta^{-1}(\mathcal{R}) = \max\{|\mathcal{R}^{-1}(x)| : x \in W\}.$$

Suppose that $|V| = |W|$. A bijective selection of $\mathcal{R}$ is called a *matching* of $\mathcal{R}$. The number of matchings of $\mathcal{R}$ will be denoted by $\mu(\mathcal{R})$.

We shall use the next two known results:

Theorem 1 (*König-Hall's Theorem*) [17] *Let $\mathcal{R} \subset V \times W$ be a relation with $|V| = |W|$. Then $\mathcal{R}$ has a matching if and only if $|\mathcal{R}(Y)| \geq |Y|$, for every subset Y of V .*

An easy consequence of König-Hall's Theorem is the following result observed by Heteyi, c.f. [17]:

Corollary 2 *Let $\mathcal{R} \subset V \times W$ be a relation with $|V| = |W|$ such that $|\mathcal{R}(Y)| \geq |Y| + 1$, for every proper subset Y of V . Then every arc $(x, y) \in \mathcal{R}$ lies in some matching of $\mathcal{R}$. In particular, $\mu(\mathcal{R}) \geq \max\{\Delta(\mathcal{R}), \Delta^{-1}(\mathcal{R})\}$.*

Some applications of Matching Theory may be found in the book of Lovász and Plummer [17]. We mention also applications of Theorem 1 and Corollary 2 by Fournier to planar tailings. [5].

Let A and B be finite subsets of a group G . The relation $\mathcal{R} = \{(x, y) \in B \times A \mid xy \notin A\}$ will be called a *Wakeford graph*. Clearly $\mathcal{R}(x) = A \cap (x^{-1}\overline{A})$, for every $x \in B$ and $\mathcal{R}(X) = A \cap (X^{-1}\overline{A})$, for every $X \subset B$.

Clearly a Wakeford pairing from B onto A is just a matching of the relation $\mathcal{R}$. Thus

$$\mu(B, A) = \mu(\mathcal{R}).$$

We shall say that the couple (B, A) is *matchable* if there is a matching from B onto A .

Notice that $xy \notin A$ if and only if $xya \notin Aa$. In particular

$$\mu(B, A) = \mu(B, Aa) \tag{1}$$

For every $x \in B$, we have $\mathcal{R}(x) = A \setminus (x^{-1}A)$. In particular,

$$|\mathcal{R}(x)| = |A \setminus (x^{-1}A)| = |xA \setminus A| = \lambda_A(x). \quad (2)$$

Since $\mathcal{R}(X) = (X^{-1}\overline{A}) \setminus \overline{A} = ((\tilde{X}^{-1})\overline{A}) \setminus \overline{A}$, Hall's condition in this graph takes the following form:

For every $X \subset B$, $|X| \leq |(\tilde{X}^{-1})\overline{A} \setminus \overline{A}|$. In particular, we have by König-Hall's Theorem and Corollary 2:

Lemma 3 *Let B be a finite subset of a group G with $1 \notin B$ and let A be a subset of G with $|A| = |B|$. Let $\mathcal{R} \subset B \times A$ be the Wakeford graph. Then*

- $\mu(B, A) \neq 0$ if and only if for every $X \subset B$, $|X| \leq |(\overline{A^{-1}\tilde{X}}) \setminus \overline{A^{-1}}|$.
- If $\mu(B, A) < \Delta(\mathcal{R})$, then there exists a proper subset $X \subset B$, with $|X| \geq |(\overline{A^{-1}\tilde{X}}) \setminus \overline{A^{-1}}|$.

2.2 The Erdős-Heilbronn averaging argument

Let G be a group, $B \subset G$ and $x \in G$. The Erdős-Heilbronn function λ is defined by the relation

$$\lambda_B(x) = |(Bx) \setminus B|.$$

Erdős and Heilbronn introduced this function in the abelian case and proved its sub-modularity [4]. Olson generalized it to the non-abelian case. We need two properties of this function:

Lemma 4 (Olson [20]) *Let B and C be nonempty subsets of a group G such that $1 \notin C$. Then,*

$$\lambda_B(x) + \lambda_B(y) \geq \lambda_B(xy). \quad (3)$$

$$\sum_{x \in C} \lambda_B(x) \geq |B|(|C| - |B| + 1). \quad (4)$$

The following lemma will be used later to show that $\Delta(\mathcal{R})$ is not small.

Lemma 5 *Let S and T be finite subsets of a group G with $1 \notin S$ and put $q = |\langle S \rangle|$. Then there is an $x \in S$ such that*

$$\lambda_T(x) \geq \min\left\{\frac{|T|(|S| + \kappa_2 - |T| + 1)}{|S| + 2\kappa_2}, \frac{|T|(q - |T| - 1)}{2q - |S| - 4}\right\},$$

where $\kappa_2 = \kappa_2(\tilde{S})$.

In particular, if $\kappa_2 = |S| = |T|$, then there is an $x \in S$ such that

$$\lambda_T(x) \geq \min\left\{\frac{|S|+1}{3}, \frac{|S|(q-|S|-1)}{2q-|S|-4}\right\}. \quad (5)$$

Proof. Put $\alpha = \max\{\lambda_T(x) : x \in S\}$.

By the definition of κ_2 , we have $|\tilde{S}^2| \geq \min(q-1, 1+|S|+\kappa_2)$. Take a subset C of $\tilde{S}^2$ such that $1 \notin C$, $S \subset C$ and $|C| = \min(q-2, |S|+\kappa_2)$

By (4) and (3),

$$\begin{aligned} |T|(|C|-|T|+1) &\leq \sum_{x \in C} \lambda_T(x) \\ &= \sum_{x \in S} \lambda_T(x) + \sum_{x \in C \setminus S} \lambda_T(x) \\ &\leq \alpha|S| + 2\alpha(|C|-|S|) \\ &= \alpha(2|C|-|S|). \end{aligned}$$

Thus $\alpha \geq \frac{|T|(|C|-|T|+1)}{2|C|-|S|}$.

Assume first that $|S| + \kappa_2 \leq q-2$. Then $|C| = |S| + \kappa_2$ and hence $\alpha \geq \frac{|T|(|S|+\kappa_2-|T|+1)}{|S|+2\kappa_2}$.

Assume now that $|S| + \kappa_2 > q-2$. Then $|C| = q-2$ and hence $\alpha \geq \frac{|T|(q-|T|-1)}{2q-|S|-4}$. ■

2.3 Isoperimetric Preliminaries

For a subset $X \subset G$, we shall write $\overline{X} = G \setminus X$ and $\tilde{X} = X \cup \{1\}$.

Let T and S be subsets of a group G with $1 \in S$. We put $T^S = G \setminus (TS)$ and $\partial_S(T) = (TS) \setminus T$. Clearly $G = T \cup T^S \cup \partial_S(T)$ is a partition. We shall write $\partial_S^-(T) = (TS^{-1}) \setminus T$. Clearly $\partial_S^-(T^S) \cap T = \emptyset$, otherwise there exist $z \in T^S$ and $y \in S$ such that $zy^{-1} = x \in T$, and hence $z = xy \in T \cup \partial_S(T)$, a contradiction. Hence

$$\partial_S^-(T^S) \subset \partial_S(T). \quad (6)$$

The last observation, used extensively in the isoperimetric method, contains a useful duality.

Recall the following result:

Theorem 6 [6, 9] *Let S be a finite subset of a group with $1 \in S$. Then there is a finite subgroup $L \neq \langle S \rangle$ generated by a subset of S such that $\kappa_1(S) = \min(|LS| - |L|, |SL| - |L|)$.*

Corollary 7 [Proposition 2.8, [8]] $\kappa_1(\tilde{S}) = |S|$, for any finite Chowla subset S .

Proof. By Theorem 6, for some finite subgroup $L \neq \langle S \rangle$, generated by a nonempty subset of S , we have $\kappa_1(\tilde{S}) = \min(|L\tilde{S}| - |L|, |\tilde{S}L| - |L|) \geq |S|$, if $|L| = 1$. Assume that $|L| \geq 2$. Since $1 \in S$, we have $\min(|LS|, |SL|) \geq 2|L|$. Thus $\kappa_1(\tilde{S}) = \min(|LS| - |L|, |SL| - |L|) \geq |L| \geq |S| + 1$, contradicting the obvious inequality $\kappa_1(\tilde{S}) \leq |S|$. ■

We need also the following more precise result:

Theorem 8 [Theorem 3.2, [8]] *Let S be a finite Chowla subset. Then $\tilde{S}$ is either a Vosper's subset or a progression.*

Let us mention that the notions of left and right progressions coincide for a subset containing 1. For this reason we shall formulate some results using translate copies of sets.

3 An inverse theorem for cofinite subsets

Inverse Theory (including the isoperimetric approach) deals only with finite sets. In this section, we derive an inverse theorem for cofinite sets.

Proposition 9 *Let S be a finite Cauchy subset of a group G with $1 \in S$. Let T be a cofinite subset of G with $|\overline{T}| \leq |\langle S \rangle| - 1$. Then $|\partial_S(T)| \geq |S| - 1$ and*

$$T^S a^{-1} \subset \langle S \rangle, \text{ for every } a \in T^S. \quad (7)$$

Assuming moreover that S is a Vosper's subset and that $|\partial_S(T)| = |S| - 1$. Then

$$(T^S)S^{-1} = T, \text{ and} \quad (8)$$

$$|(T^S)S^{-1}| = |T^S| + |S| - 1. \quad (9)$$

Proof. Put $Z = T^S$ and $H = \langle S \rangle$. Take a left-decomposition $Z = Z_1 \cup \dots \cup Z_j$ modulo H , where $Z_i = z_i H \cap Z$, for some $z_i \in Z$. By (6),

$$Z_i S^{-1} \subset Z_i \cup \partial_S^-(Z_i) \subset Z_i \cup \partial_S(Z_i) \subset \overline{T},$$

and hence $|Z_i S^{-1}| \leq |\overline{T}| \leq |H| - 1$. By the definition of κ_1 , we have $|Z_i S^{-1}| = |(z_i^{-1} Z_i) S^{-1}| \geq \min(|H|, |Z_i| + |S| - 1) = |Z_i| + |S| - 1$. Therefore $|Z S^{-1}| = \sum_{1 \leq i \leq j} |Z_i S^{-1}| \geq |Z| + j(|S| - 1)$. Thus we have by (6), $j(|S| - 1) \leq |\partial_S^-(Z)| \leq |\partial_S(T)| \leq |S| - 1$. Thus $j = 1$, proving (7). We have also

$$|S| - 1 \leq |\partial_S^-(T^S)| \leq |\partial_S(T)| \quad (10)$$

Assume now that S is a Vosper's subset and that $|\partial_S(T)| \leq |S| - 1$. By (10), $|S| - 1 \leq |\partial_S^-(T^S)| \leq |\partial_S(T)| \leq |S| - 1$.

In particular, $|(T^S)S^{-1}| = |T^S| + |\partial_S^-(T^S)| = |T^S| + |S| - 1$, proving (9).

By (10) and (6),

$$(T^S)S^{-1} = T^S \cup \partial_S^-(T^S) = T^S \cup \partial_S(T) = \overline{T}.$$

■

We can now show that Chowla subsets behave nicely with respect to matchability.

Theorem 10 *Let B be a finite Chowla subset of a group G . If A be a subset of G with $|A| = |B|$, then $\mu(B, A) \neq 0$.*

Proof. Let X be an arbitrary subset of B . Put $H = \langle X \rangle$ and $\tilde{X} = U$. Clearly U is a Chowla subset. By Corollary 7, U is a Cauchy subset. Put $V = (\overline{A})^{-1}$ and $W = V^U$.

By Proposition 9, $|\overline{A^{-1}}\tilde{X} \setminus \overline{A^{-1}}| = |\partial_U(V)| \geq |U| - 1 = |X|$. By Lemma 3, $\mu(B, A) \neq 0$. ■

In particular,

Corollary 11 *Let B be a finite subset of a group G with $1 \notin B$ and let A be a subset of G with $|A| = |B|$.*

If $|B| < p(G)$, then $\mu(B, A) \neq 0$.

Applied to groups with a prime order, the last result reduces to a result of Losonczy [16]. Also in the case of torsion free groups, it reduces to a result of Eliahou-Lecouvey [1] generalizing to the non-abelian case results of Wakeford [24] and Losonczy [16].

4 Distinct matchings

The Vosper's property implies a bound the number of distinct matchings. We shall illustrate this relation in the easier case of Chowla subsets.

Theorem 12 *Let B be a finite Chowla subset of a group G let A be a subset of G with $|A| = |B|$. Let $\mathcal{R} \subset B \times A$ be the Wakeford graph. Then one of the following holds:*

- (i) $\mu(B, A) \geq \max\{\Delta(\mathcal{R}), \Delta^{-1}(\mathcal{R})\}$.
- (ii) For some $a \in A$, $|Aa^{-1} \cap B| = |B| - 1$.
- (iii) For some $a \in A$, Aa^{-1} is a progression.

Proof. Suppose that (i) is not satisfied. By Lemma 3, for some proper subset X of B , we have $|(\overline{A^{-1}\tilde{X}}) \setminus \overline{A^{-1}}| = |\partial_U(V)| \leq |U| - 1 = |X|$, where $\tilde{X} = U$ and $V = \overline{A^{-1}}$.

Put $H = \langle X \rangle$ and $W = V^U$. Take $w \in W$.

By Proposition 9, $|\partial_U(V)| = |U| - 1 = |X|$. By (8),

$$A^{-1} = WU^{-1}.$$

Assume first that $|W| = 1$, and hence $W = \{w\}$. Notice that $w \in A^{-1}$. We have clearly $Aw = U$. Thus $|A| = |U| = |X| + 1$ and hence $|X| = |B| - 1$. Therefore $|Aw \cap B| = |X| = |B| - 1$, and (ii) holds. Assume $|W| \geq 2$. By Theorem 8, we have one of the two possibilities:

- U is an r -progression, for some r .

By (9), $|UW^{-1}| = |W| + |U| - 1 = |A| \leq |H| - 1$. It follows that $W^{-1}w$ is a progression and hence $UW^{-1} = Aw$ is a progression. Thus (iii) holds.

- U is a Vosper subset.

Then clearly $|X| \geq |U| - 1 \geq 2$, (since a subset of size 2 is not a Vosper's subset). Choose $r \in X$.

By (9), $|UW^{-1}| = |W| + |U| - 1 = |A| \leq |H| - 1$. Since U is a Vosper's subset of H , we must have $|A| = |UW^{-1}| = |H| - 1$. By (7), $A \subset Hw$, and hence $|(Aw^{-1}) \cap H| = |H| - 1$. But $|\langle r \rangle| \geq |A| + 1 = |H|$, and hence $H = \langle r \rangle$. Since H is cyclic, any subset of H with size $|H| - 1$ is a progression. Thus Aw^{-1} is a progression and (iii) holds

■

Corollary 13 *Let B be a Chowla subset with $q = |\langle B \rangle|$. Then one of the following conditions holds:*

$$(i) \mu(B, B) \geq \min\left\{\frac{|B|+1}{3}, \frac{|B|(q-|B|-1)}{2q-|B|-4}\right\}.$$

(ii) *For some $a \in B$, Ba^{-1} is a progression.*

Proof. Let $\mathcal{R} \subset B \times A$ be the Wakeford graph. Suppose that (i) and (ii) are false. By (5) and (2), $\Delta(\mathcal{R}) \geq \min\left\{\frac{|B|+1}{3}, \frac{|B|(q-|B|-1)}{2q-|B|-4}\right\} > \mu(\mathcal{R})$. By Theorem 12, for some $a \in B$, $|Ba \cap B| = |B| - 1$.

Since $a \neq 1$, we have $|B\{1, a\}| = |B| + 1$. Since $|B|$ is less than the order of a , B is an a -progression. Thus Ba^{-1} is a progression, a contradiction. ■

References

- [1] S. Eliahou and C. Lecouvey, Matchings in arbitrary groups, *Advances in Applied Mathematics* 40 (2008) 219224.
- [2] C. K. Fan and J. Losonczy, Matchings and canonical forms in symmetric tensors, *Adv. Math.* 117 (1996), 228-238.
- [3] P. Hall, On representatives of subsets, *J. London Math. Soc.* 10 (1935), 26-30.
- [4] P. Erdős and H. Heilbronn, On the Addition of residue classes mod p , *Acta Arith.* 9 (1964), 149-159.

- [5] J. C. Fournier, Combinatorics of perfect matchings in plane bipartite graphs and application to tilings. *Theoret. Comput. Sci.* 303 (2003), no. 2-3,
- [6] Y.O. Hamidoune, On the connectivity of Cayley digraphs, *Europ. J. Combinatorics*, 5 (1984), 309-312.
- [7] Y.O. Hamidoune, On subsets with a small sum in abelian groups I: The Vosper property, *Europ. J. of Combinatorics* 18 (1997), 541-556.
- [8] Y.O. Hamidoune, An isoperimetric method in additive theory. *J. Algebra* 179 (1996), no. 2, 622-630.
- [9] Y.O. Hamidoune, On small subset product in a group. Structure Theory of set-addition, *Astérisque* no. 258(1999), xiv-xv, 281-308.
- [10] Y.O. Hamidoune, Some results in Additive number Theory I: The critical pair Theory, *Acta Arith.* 96, no. 2(2000), 97-119.
- [11] Y.O. Hamidoune, Hyper-atoms and the Kemperman's critical pair Theory, Preprint, septembre 2007.
- [12] G. Károlyi, Cauchy-Davenport theorem in group extensions, *Enseign. Math.* (2) 51 (2005), no. 3-4, 239-254.
- [13] J.H.B. Kempermann, On complexes in a semigroup, *Nederl. Akad. Wetensch. Proc. Ser. A.* 59 *Indag. Math.* 18 (1956), 247-254.
- [14] M. Kneser, Abschätzung der asymptotischen Dichte von Summenmengen. *Math. Z.* 58, (1953). 459-484.
- [15] M. Kneser, Summenmengen in lokalkompakten abelschen Gruppen. *Math. Zeit.* 66 (1956), 88-110.
- [16] J. Losonczy, On matchings in groups, *Adv. in Appl. Math.* 20 (1998) 385-391.
- [17] L. Lovász, M.D. Plummer, Matching theory, *Ann. Discrete Math.* 29 (1986) 137.
- [18] M. B. Nathanson, *Additive Number Theory. Inverse problems and the geometry of sumsets*, Grad. Texts in Math. 165, Springer, 1996.
- [19] J. E. Olson, An addition theorem mod p , *J. Comb. Theory* 5 (1968), 45-52.

- [20] J.E. Olson, Sums of sets of group elements, *Acta Arith.* 28 (1975/76), no. 2, 147–156.
- [21] J.E. Olson, On the sum of two sets in a group, *J. Number Theory* 18 (1984), 110–120.
- [22] P. Scherk, L.Moser, Advanced Problems and Solutions: Solutions: 4466, *Amer. Math. Monthly* 62 (1955), no. 1, 46–47.
- [23] T. Tao and V.H. Vu, *Additive Combinatorics*, Cambridge Studies in Advanced Mathematics 105 (2006), Cambridge Press University.
- [24] E.K. Wakeford, On canonical forms, *Proc. London Math. Soc.* 1818 (1918/1919), 403–410.